



Cluster	Product (Group)	Registration #	Application placement	Formulation type	WALES mixing order	Maximum applications*	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
Biologicals														
Biostimulant	 Amatis™	2017126F	Drench or soil	L	S	NA	1.4 L/acre (3.5 L/ha)	None	14 to 21	0	0			Helps to revitalize and improve the rhizosphere around plant roots, promoting root growth allowing for improved yield and uniformity.
	 Megafol™	2016013D	Foliar	L	S	NA	1.0 L/acre (2.5 L/ha)	None	10 to 15	0	0			Helps plants respond to abiotic stress including physical injury, heat, cold and drought. Consider as a regular part of your spray program to assist with stress. In case of physical injury apply as soon as possible after.
Fungicides														
Phytophthora	 Orondis™ Gold DC (Group 4 + 49)	35176	Soil	EC	E	3	1.21 L/acre (3 L/ha)	None	42	12	14	Phytophthora root rot, Pythium damping off		Maximum of 3 applications of group 4-containing products permitted per season (Ridomil® Gold, Orondis® Gold).
	 Revus™ (Group 40)	29074	Foliar or soil	SC	L	4	0.24 L/acre (0.58 L/ha)	NIS 0.25% v/v	7	12	3	Phytophthora blight and root rot		
	 RidomilGold® (Group 4)	28474	Soil	SL	S	1	0.26 L/acre (0.65 L/ha)	None	NA	12	9	Phytophthora root rot, Pythium damping off		Maximum of 3 applications of group 4-containing products permitted per season (Ridomil Gold, Orondis Gold).
	 RidomilGold™ 1G (Group 4)	26612	Soil	G	NA	3	12.6 kg/acre (31.25 kg/ha)	None	42	12	9	Phytophthora root rot		Maximum of 3 applications of group 4-containing products permitted per season (Ridomil Gold, Orondis Gold).
Root rots	 Allegro™ (Group 29)	27517	Soil	SC	L	4	0.49 L/acre (1.2 L/ha)	None	14	24	30	Rhizoctonia root rot		
	 CaptanL™ by syngenta. (Group M04)	35343	Foliar	SC	L	8	1.68 L/ac (4.15L/ha)	None	7	7 days hand set irrigation; 12 hours other activities	20	Rhizoctonia root rot, Pythium root rot and damping-off, Phytophthora root rot, grey mould	Cylindrocarpon root rot	The application rate should be based on 4.15 L product per hectare applied in the appropriate volume of carrier to achieve thorough coverage of all plant surfaces.
	 Quadris™ (Group 11)	26153	Soil	SC	L	1	0.45 L/acre (1.12 L/ha)	None	NA	12	90	Rhizoctonia root rot		
	 Switch™ (Group 9 + 12)	28189	Soil	WG	W	2	0.31 to 0.4 kg/acre (775 to 975 g/ha)	None	14 to 21	12	7	Rhizoctonia root rot	Cylindrocarpon	Rate differs by target.
Leaf blights	 Allegro™ (Group 29)	27517	Foliar	SC	L	4	0.49 L/acre (1.2 L/ha)	None	7 to 14	24	30	Alternaria, Botrytis		
	 Aprovia™ Top (Group 3 + 7)	31526	Foliar	EC	E	4	0.26 to 0.39 L/acre (0.64 to 0.97 L/ha)	None	10 to 14	12	14	Alternaria		Do not exceed 300 g of benzovindiflupyr (Solatenol®) or 500 g of difenoconazole per hectare per season. Aprovia®Top fungicide contains 78 g of benzovindiflupyr and 117 g of difenoconazole per litre.
	 Miravis™ Duo (Group 3 + 7)	33206	Foliar	SC	L	4	0.4 L/acre (1 L/ha)	None	7 to 14	12	7	Alternaria	Botrytis	Do not exceed 300 g of pydiflumetofen (ADEPIDYN®) or 500 g of difenoconazole per hectare per season. Miravis® Duo fungicide contains 75 g of pydiflumetofen and 125 g of difenoconazole per litre.
	 Switch™ (Group 9 + 12)	28189	Foliar	WG	W	2	0.31 to 0.4 kg/acre (775 to 975 g/ha)	None	7 to 10	12	7	Botrytis, Alternaria		Rate differs by target. <i>Botrytis cinerea</i> populations with reduced sensitivity to Group 9 fungicides have been observed in some regions of Canada.
	 Bravo™ZN Bravo™ZNC (Group M)	28900 33515	Foliar	SC	L	3	0.97 to 1.94 L/acre (2.4 to 4.8 L/ha)	None	10	12	14	Alternaria, Botrytis		Two applications in-season and one application in the fall.

Please check with your exporter when using any Syngenta product on ginseng destined for use outside of Canada. At time of print, Maximum Residue Limits (MRLs) had been established for the active ingredients benzovindiflupyr and fluazinam for export markets in the United States, pydiflumetofen, fludioxonil and cyprodinil for export markets in the United States and Codex, oxathiapiprolin for export markets in Codex, the United States and China, azoxystrobin for export markets in Codex, the United States, China and Hong Kong, mandipropamid for export markets in the United States and China, and chlorothalonil and metalaxyl-m for export markets in the United States, Codex, China and Hong Kong. Please see the Foodchain ID Regulatory Limits by BCGlobal database at <https://bcglobal.bryantchristie.com/db#/pesticides/query> for a complete list of MRLs. Should you need additional information on export market MRLs, please consult with Syngenta to receive the most up-to-date information.

REI = Re-entry interval


PHI = Pre-harvest interval

Cluster	Product (Group)	Registration #	Application placement	Formulation type	WALES mixing order	Maximum applications*	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
Herbicide														
Grasses	 Venture [®] L (Group 1)	21209	Post-emergent	EC	E	3	0.24 to 0.81 L/acre (0.6 to 2 L/ha)	None	NA	12	365	Annual grasses		Application rate differs by target.

 Application timing

				
Pre-emergence/bud swell	Emergence/hooking	Vegetative	Root bulking	Senescence
				
				
				
				
				
				
				
				
				
				
				
				

Resources ripe for the picking



Access crop protection insights to grow your most marketable crop.



*For complete information regarding rates, see product label. Rate information is based on product labels. If a product label contains rate ranges, then this quick reference guide will list either: (i) the complete rate range; (ii) the most common rate range; or (iii) the most commonly used rate.

Always refer to the label to confirm crop/crop group registration and application information, as well as buffer zones, PPE, etc., before application.

All information is current at time of publication and is subject to change without notice. For more information, contact our Customer Interaction Centre at 1-877-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

Always read and follow label directions. ADEPIDYN[®], Aprovia[®], Bravo[®], Miravis[®], Orondis[®], Quadris[®], Revus[®], Ridomil Gold[®], Scholar[®], Solatenol[®], Switch[®], Venture[®] and the Syngenta logo are trademarks of a Syngenta Group Company. Amatis[™] and Megafol[™] are trademarks of VALAGRO S.p.A., a Syngenta Group Company. Allegro[®] is a trademark of ISK Biosciences Corporation. Used under license. Other trademarks are the property of their respective owners. © 2025 Syngenta

250929