
luster	Product (Group)	Registration #	Crop (crop group)	Application placement	Formulation type	WALES mixing order	Maximum applications*	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
Ö	Biostimulant														
nulant	O Amatis	2017126F	All	Drench or soil	L	S	NA	1.4 L/acre (3.5 L/ha)	None	14 to 21	0	0			Helps to revitalize and improve the rhizosphere around plant roots, promoting root growth allowing for improved yield and uniformity.
Biostin	Megafol [*]	2016013D	All	Foliar	L	S	NA	1.0 L/acre (2.5 L/ha)	None	10 to 15	0	0			Helps plants respond to abiotic stress including: physical injury, heat, cold and drought. In case of physical injury apply as soon as possible after.
	Fungicide														
Clubroot	Allegro ° (Group 29)	27517	Brassica leafy (5)	Drench or banded	SC	L	1	1.2 L/acre 50 mL/100 L (2.9 L/ha)	None	NA	24	5A crops: 65 5B crops: 30	Clubroot		
	Bravo ZN	28900	Cabbage	Foliar	SC	L	2	1 to 1.9 L/acre (2.5 to 4.8 L/ha)	None	7 to 10	2 days for scouting and thinning,1 day for hand weeding,12 hours for all other activities	7	Alternaria, downy mildew		
	(Group M)		Broccoli, Brussels sprouts, cauliflower	Foliar	SC	L	1	1 to 1.9 L/acre (2.5 to 4.8 L/ha)	None	NA	5 days for topping and scouting, 12 hours for all other activities	7	Alternaria, downy mildew		
	×		Brassica leafy (4-13B)	Foliar	SC	L	3	0.3 to 0.4 L/acre (0.73 to 1 L/ha)	NIS 0.125% v/v	7 to 10	4 days for weeding, 12 hours for all other activities	3	Alternaria, powdery mildew		
Leaf spots	XX Miravis° Duo (Group 3 + 7)	33206	Brassica head and stem (5-13)	Foliar	SC	L	3	0.3 to 0.4 L/acre (0.73 to 1 L/ha)	NIS 0.125% v/v	7 to 10	5 days for hand harvesting, 4 days for weeding, 3 days for scouting, topping, tying and training, 12 hours for all other activities	5	Alternaria, powdery mildew		
	Quadris ° (Group 11)	26153	Cabbage	Foliar	SC	L	3	0.45 L/acre (1.12 L/ha)	NIS 0.125% v/v	7 to 14	12	1	Alternaria		
	Quadris Top° (Group 3 + 11)	30518	Brassica leafy (5)	Foliar	SC	L	4	0.29 to 0.4 L/acre (0.71 to 1 L/ha)	NIS 0.125% v/v	10 to 14	3 days for hand harvesting, 1 day for scouting, 12 hours for all other activities	3	Alternaria, powdery mildew		
	(Group 9 + 12)	28189	Brassica leafy (4-13B) brassica head and stem (5-13)	Foliar	WG	W	3	313 to 395 g/acre (775 to 975 g/ha)	NIS 0.125% v/v	7 to 10	12	7	Powdery mildew	Alternaria	Target disease differs by crop.
nildew	Orondis'Ultra (Group 40 + 49)	32805	Brassica leafy greens (4-13B) and brassica head and stem (5-13)	Foliar	SC	L	4	0.24 L/acre (0.6 L/ha)	NIS 0.125% v/v	7 to 10	12	1	Downy mildew		
Downy n	Revus [®] (Group 40)	29074	Brassica leafy greens (4-13B), brassica head and stem (5-13) and kohlrabi	Foliar	SC	L	4	0.16 to 0.24 L/acre (0.4 to 0.6 L/ha)	NIS 0.125% v/v	7 to 10	12	1	Downy mildew		
	Insecticide														
Chewing and sucking pests	Actara 240SC (Group 4)	28407	Brassica leafy (5)	In-furrow or banded	SC	L	1	0.15 L/ac (0.375 L/ha)	None	NA	12	NA	Aphids	Flea beetles	
	Herbicides														
Broadleaves and grasses	Dual I Magnum ® (Group 15)	25729	Broccoli, cabbage and cauliflower	Pre-emergent or post- emergent	EC	E	1	0.5 to 0.71 L/acre (1.25 to 1.75 L/ha)	None	NA	12	NA	Nightshade, annual grasses	Redroot pigweed	Transplanted crops only.
Grasses	Venture L (Group 1)	21209	Broccoli, Brussels sprouts, cabbage and cauliflower	Post-emergent	EC	Е	1	0.32 to 0.81 L/acre (0.8 to 2 L/ha)	None	NA	12	40	Annual grasses		Application rate differs by target.







Brassica application timing



^{*}For complete information regarding rates, see product label. Rate information is based on product labels. If a product label contains rate ranges, then this quick reference guide will list either: (i) the complete rate range; (ii) the most common rate range; or (iii) the

All information is current at time of publication and is subject to change without notice. For more information, contact our Customer Interaction Centre at 1-87-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

Always read and follow label directions. Actara®, Bravo®, Dual II Magnum®, Miravis®, Orondis®, Quadris®, Quadris®, Quadris®, Amatis™ and Megafol™ are trademarks of VALAGRO S.p.A., a Syngenta Group Company. Allegro® is a trademark of ISK Biosciences Corporation. © 2025 Syngenta.

