
Cluster	Product (Group)	Registration #	Application placement	mixing		Maximum applications	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
O	Fungicides													
ts	RidomilGold® (Group 4)	28474	Soil and drip	SL	S	1	37 mL/100 m row	None	NA	12	80	Phytophthora		
Root rots	Orondis Gold DC Orondis Gold (Group 4 + 49)	35176 33508	Soil and drip	EC	E	2	1.85 L/acre (4.57 L/ha)	None	30	12	1	Phytophthora		For effective disease control, ensure that the product solution thoroughly wets the target root zone.
	<b>Allegro</b> * (Group 29)	27517	Foliar	SC	L	2	0.91 L/acre (2.24 L/ha)	None	7 to 10	24	30	Botrytis grey mould	Mummyberry, Phomopsis fruit rot, fruit anthracnose	
	Bravo ZNC (Group M)	28900 33515	Foliar	SC	L	2	2.9 L/acre (7.2 L/ha)	None	NA	72 for handset irrigation, 12 for all other activities	54	Anthracnose, Alternaria, Phomopsis		
	(TinspireSuper (Group 3 + 9)	30827	Foliar	EW	Е	4	0.23 to 0.6 L/acre (0.59 to 1.48 L/ha)	None	10 to 21	12	1	Alternaria, anthracnose, Botrytis grey mould, mummyberry, Monilinia	Rust	Application rate differs by target.
Fruit rots	<b>X Miravis Neo</b> (Group 3 + 7 + 11)	33391	Foliar	SE	L	2	0.3 L/acre (0.75 L/ha)	None	7	12	30	Mummyberry, Monilinia blight, anthracnose		Do not make more than 2 applications of any ADEPIDYN® containing product (e.g. Miravis® Neo or Miravis® Prime) on blueberries per year.
	Quilt° (Group 3 + 11)	28328	Foliar	SE	L	2	0.4 L/acre (1 L/ha)	None	7 to 14	12	30	Mummyberry, Monilinia blight, anthracnose, rust	Septoria leaf spot, Valdensinisa leaf spot	
	* <b>X Miravis°Prime</b> (Group 7 + 12)	33207	Foliar	SC	L	2	0.32 to 0.4 L/acre (0.8 to 1.0 L/ha)	None	7 to 10	12	1	Botrytis fruit rot		Do not make more than 2 applications of any ADEPIDYN® containing product (e.g. Miravis® Neo or Miravis® Prime) on blueberries per year.  Do not apply more than 552 g of fludioxinil per ha per season.
	<b>Switch</b> * (Group 9 + 12)	28189	Foliar	WG	W	3	314 to 395 g/acre (775 to 975 g/ha)	None	7 to 10	12	1	Botrytis grey mould, Monilinia, mummyberry, anthracnose		Do not apply more than 552 g of fludioxinil per ha per season.
	Herbicides													
Broadleaves	<b>Callisto®</b> (Group 27)	27833	Banded	SC	L	1	121 mL/acre (0.3 L/ha)	Post- emergent: Agral® 90 0.2% v/v	NA	12	NA	Pre-emergent: Lamb's-quarters, redroot pigweed, velvetleaf Post-emergent: Nightshade, velvetleaf, redroot pigweed	Common ragweed	
Grasses and broadleaves	<b>Dual I Magnum</b> ° (Group 15)	25729	Pre- emergent	EC	E	1	0.51 to 0.71 L/acre (1.25 to 1.75 L/ha)	None	NA	12	30	Nightshade (American and Eastern black), crabgrass (smooth, hairy), barnyard grass, fall panicum, foxtail (green, yellow, giant), old witchgrass	Redroot pigweed	
Grasses	Venture L (Group 1)	21209	Banded	EC	E	1	0.24 to 0.81 L/acre (0.6 to 2 L/ha)	None	NA	12	15	Annual and perennial grasses		Application rate differs by target.



REI = Re-entry interval

PHI = Pre-harvest interval



## Resources ripe for the picking




Access crop protection insights to grow your most marketable crop.



## 2025 Highbush blueberry quick reference guide

Cluster	Product (Group)	Registration #	Application placement	Formulation type	WALES mixing order	Maximum applications	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Comments
ပ	Biologicals											
Biostimulant		2016013D	Foliar	L	S	1	1.0 L/acre (2.5 L/ha)	None	10 to 15	0	0	Helps plants respond to unexpected abiotic stress including: physical injury, heat, cold, excess moisture, drought. In case of physical injury, apply as soon as possible after.
Biost		2017126F	Drench or soil	L	S	NA	2.0 L/acre (5.0 L/ha)	None	14 to 21	0	0	Helps to revitalize and improve the rhizosphere around plant roots, promoting root growth allowing for improved yield and uniformity.

## Application timing



Please check with your packer or exporter when using Allegro®, Bravo® ZNC/ZN, Dual® II Magnum, Miravis® Neo, Miravis® Prime, Orondis® Gold, Quilt®, Ridomil® Gold and Venture® L on berries destined for use outside of Canada. At time of print, Maximum Residue Limits (MRLs) had been established for the active ingredients fluazifop-p-butyl, oxathiapiprolin and S-metolachlor for export markets in the United States, and azoxystrobin, chlorothalonil, fluazinam, metalaxyl-M,

Please see the Foodchain ID Regulatory Limits by BCGlobal database at https://bcglobal.bryantchristie.com/db#/ pesticides/query for a complete list of MRLs. Should you need additional information on export market MRLs, please consult with Syngenta to receive the most up-to-date information.

pydiflumetofen and propiconazole for export markets in the United States and Japan.

All information is current at time of publication and is subject to change without notice.

For more information, contact our Customer Interaction Centre at 1-87-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

Always read and follow label directions. Miravis® Neo refers to Miravis® Neo 300SE fungicide. ADEPIDYN®, Agral®, Aprovia®, Bravo®, Callisto®, Dual II Magnum®, Inspire Super®, Miravis®, Orondis®, Quilt®, Ridomil Gold®, Switch®, Venture® and the Syngenta logo are trademarks of a Syngenta Group Company. Amatis™ and Megafol® are trademarks of VALAGRO S.p.A., a Syngenta Group Company. Allegro® is a trademark of ISK Biosciences Corporation. © 2024 Syngenta

240814

