









































Cluster	Product (Group)	Registration #	Application placement	Formulation type	WALES mixing order	Maximum applications*	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
Biologicals														
Biostimulant	 <b>Amatis™</b>	2017126F	Drench or soil	L	S	NA	2 L/acre (5 L/ha)	None	14 to 21	0	0			Amatis™ helps to revitalize and improve the rhizosphere around plant roots, promoting root growth allowing for improved yield and uniformity.
	 <b>Megafol™</b>	2016013D	Foliar	L	S	NA	1.0 L/acre (2.5 L/ha)	None	10 to 15	0	0			Helps plants respond to abiotic stress including: physical injury, heat, cold, and drought. In case of physical injury apply as soon as possible after.
Fungicides														
Root rots	 <b>Quadris®</b> (Group 11)	26153	Banded and drip	SC	L	2	0.45 L/acre (1.1 L/ha)	None	NA	12	1		Black root rot	
	 <b>Orondis® Gold DC</b> (Group 4 + 49)	35176	Banded and drip	EC	E	2	1.85 L/acre (4.57 L/ha)	None	7	12	1	Red stele, crown rot, Phytophthora root rot, red stele root rot	Phytophthora crown rot	Do not reduce rate for banded applications.
	 <b>Ridomil® Gold</b> (Group 4)	28474	Foliar and banded	SL	S	2	Eastern Canada: 0.4 L/acre (1 L/ha) Western Canada: 20 mL/100 m	None	30	12	NA	Red stele		New plantings: Apply to the soil surface in a one-metre wide band centered over the row, as a post planting soil drench. Make a second application at the same rate in the fall prior to freeze-up Established plantings: Apply to the soil surface in a one-metre wide band centred over the row, in the fall prior to freeze-up.
	 <b>Scholar®</b> (Group 12)	29528	Banded and drip	SC	L	2	0.49 L/acre (1.2 L/ha)	None	NA	12	1		Black root rot	
Fruit rots	 <b>Bravo® ZN</b> (Group M)	28900	Foliar	SC	L	3	1.4 L/acre (3.5 L/ha)	None	10 to 12	12	30	Botrytis		Apply once in the fall. Apply two pre-bloom sprays the following spring, one when new growth appears and again 10-12 days later.
	 <b>Bravo® ZNC</b> (Group M)	33515												
	 <b>Captan® L</b> (Group M04)	35343	Foliar	SC	L	6	2.34 L/ac (5.8L/ha)	None	7	9 days for hand set irrigation, 12 hours all other activities	8	Gray-mold (Botrytis) and leaf spot (Mycosphaerella fragariae)		Use 1,000 litres water per ha.
	 <b>Inspire® Super</b> (Group 3 + 9)	30827	Foliar	EW	L	3	470 to 597 mL/acre (1161 to 1475 mL/ha)	None	7 to 10	12	1	Botrytis	Anthracnose	<i>Botrytis cinerea</i> populations with reduced sensitivity to Group 9 fungicides have been observed in some regions of Canada.
	 <b>Miravis® Prime</b> (Group 7 + 12)	33207	Foliar	SC	L	2	0.4 L/acre (1 L/ha)	None	7 to 10	12	1	Botrytis, anthracnose		
	 <b>Quadris® Top</b> (Group 3 + 11)	30518	Foliar	SC	L	3	0.35 to 0.4 L/acre (0.87 to 1.0 L/ha)	None	7 to 14	12	1	Anthracnose, powdery mildew		Resistance to the QoI fungicides has been reported; performance may be impacted.
	 <b>Switch®</b> (Group 9 + 12)	28189	Foliar	WG	W	3	314 to 395 g/acre (775 to 975 g/ha)	None	7 to 10	12	1	Botrytis, anthracnose	Powdery mildew	<i>Botrytis cinerea</i> populations with reduced sensitivity to Group 9 fungicides have been observed in some regions of Canada.
Insecticides														
Mites	 <b>Agri-Mek® SC</b> (Group 6)	31607	Foliar	SC	L	2	91 mL/acre (225 mL/ha)	NIS 0.25% v/v		12	see comment	Cyclamen mite, two-spotted spider mite, McDaniel spider mite		Pre-harvest application is 3 days. Post-harvest application is 10 months.
Broad spectrum	 <b>Matador®</b> (Group 3)	24984	Foliar	EC	E	3	42 mL/acre (104 mL/ha)	None	7	12	7	Bud clipper weevil, meadow spittle bug, tarnished plant bug	Black vine weevil (adults)	Apply in the early morning before temperatures rise, or in the evening after the heat of the day.
Herbicides														
Grasses and broadleaves	 <b>Dual® II Magnum</b> (Group 15)	25729	Pre-plant incorporated and pre-emergent	EC	E	1	0.51 to 0.71 L/acre (1.25 to 1.75 L/ha)	None	NA	12	30	Annual broadleaves and grasses, nutsedge		Do not use on Joliette. For band application, use proportionately less material. See label for application specifics.
	 <b>Venture® L</b> (Group 1)	21209	Post-emergent	EC	E	1	0.24 to 0.81 L/acre (0.6 to 2 L/ha)	None	NA	12	30	Annual and perennial grasses		Allow a buffer of 5-7 days prior to, or following, a broadleaf herbicide application. See label for application specifics.
Broadleaves	 <b>Reflex®</b> (Group 14)	24779	Dormancy	EC	E	1	0.4 L/acre (1 L/ha)	Turbocharge® 0.25% v/v or NIS 0.1% v/v	NA	12	77	Pigweed, ragweed, lamb's-quarters, wild mustard, lady's thumb, eastern black nightshade, cocklebur		For use in Eastern Canada and British Columbia only. Perennial strawberries only. See label for application specifics.

REI = Re-entry interval      PHI = Pre-harvest interval

🍓 Application timing

						
Dormant	Vegetative	Early bloom/First hand	Full bloom	Green fruit	Pre-harvest	Harvest
						
						
						
						
						
						
						
						
						
						
						
						
						
						

🍓 Strawberry quick reference guide

### Resources ripe for the picking



Access crop protection insights to grow your most marketable crop.



\*For complete information regarding rates, see product label. Rate information is based on product labels. If a product label contains rate ranges, then this quick reference guide will list either: (i) the complete rate range; (ii) the most common rate range; or (iii) the most commonly used rate. Always refer to the label to confirm crop/crop group registration and application information, as well as buffer zones, PPE, etc., before application.

All information is current at time of publication and is subject to change without notice. For more information, contact our Customer Interaction Centre at 1-877-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

**Always read and follow label directions.** Agri-Mek®, Bravo®, Dual II Magnum®, Inspire Super®, Matador®, Miravis®, Orondis®, Quadris®, Quadris Top®, Reflex®, Ridomil Gold®, Scholar®, Switch®, Turbocharge®, Venture® and the Syngenta logo are trademarks of a Syngenta Group Company. Amatis™ and Megafol™ are trademarks of VALAGRO S.p.A., a Syngenta Group Company. Used under license. All other trademarks are the property of their respective owners. © 2025 Syngenta

251110