	<b>4</b> 2
-	9
٠	$\equiv$
	ಠ
	<b>(</b> )
	$\ddot{\circ}$
	$\subseteq$
Ų	<u>H</u>
	<u>U</u>
	iick reterence
	$\overline{\mathcal{O}}$
	$\supseteq$
	O
	$\geq$
	(I)
	Ŏ
	Φ
-	=
-	
-	
	bush blueberry quick reterence guide
	9
	$\leq$
	*

Cluster	Product (Group)	Registration #	Application placement	Formulation type	WALES mixing order	Maximum applications	Application rate*	Adjuvant	Interval (days)	REI (hours)	PHI (days)	Control of	Suppression of	Comments
ਠ	Biologicals													
Biostimulant	<b>○</b> Megafol <sup>-</sup>	2016013D	Foliar	L	S	1	1.0 L/ac (2.5 L/ha)	None	10 to 15	0	0			Helps plants respond to abiotic stress including: physical injury, heat, cold, and drought. In case of physical injury, apply as soon as possible after.
	Fungicides													
Leaf spots and rust	· <b>·· Aprovia</b> (Group 7)	31981	Foliar	EC	E	2	0.2 to 0.3 L/acre (0.5 to 0.75 L/ha)	NIS 0.2% v/v	10 to 14	12	365	Rust	Valdensinia	Crop safety has been found to be acceptable; however, not all blueberry clones have been tested under all conditions. It is recommended to test Aprovia® on a representative portion of the crop and in local conditions to ensure that a phytotoxic response will not occur. Application rate differs by target.
Leaf spo	Bravo <sup>°</sup> ZN Bravo <sup>°</sup> ZNC (Group M)	28900 33515	Foliar	SC	L	2	2.9 L/acre (7.2 L/ha)	None	42	12	54	Phomopsis, Valdensinia, Septoria, rust		
	<b>XX Miravis<sup>®</sup>Neo</b> (Group 3 + 7 + 11)	33391	Foliar	SE	L	2	0.3 L/acre (0.75 L/ha)	None	10 to 14	12	365	Rust	Septoria, Valdensinia	
	Quilt° (Group 3 + 11)	28328	Foliar	SE	L	2 3 (anthracnose only)	0.4 L/acre (1 L/ha)	None	7 to 14	12	30	Rust, mummyberry, Monilinia blight, anthracnose	Septoria, Valdensinia	
Fruit rots	Allegro* (Group 29)	27517	Foliar	SC	L	2 (fruit rots) 4 (Valdensinia)	0.59 to 0.91 L/acre (1.46 to 2.24 L/ha) 0.16 to 0.32 L/acre (0.4 to 0.8 L/ha)	None	7 to 10	24	30	Botrytis grey mould	Mummyberry, Phomopsis fruit rot, fruit anthracnose, Valdensinia	Application rate differs by target.  Apply at 1.46 to 2.24 L/ha for mummyberry, fruit anthracnose, and Botrytis and 2.24 L/ha for Phomopsis.  For Valdensinia, apply at 0.4 to 0.8 L/ha.
Fru	<b>(<sup>*</sup> Inspir∈Super</b> ° (Group 3 + 9)	30827	Foliar	EW	E	2 (Alternaria, Botrytis, Monilinia) 3 (anthracnose) 4 (mummyberry)	0.23 to 0.6 L/acre (0.56 to 1.48 L/ha)	None	10 to 21	12	1	Alternaria, anthracnose, Botrytis grey mould, Monilinia, mummyberry	Rust	Application rate differs by target.  Botrytis cinerea populations with reduced sensitivity to Group 9 fungicides have been observed in some regions of Canada.
	<b>Switch</b> ° (Group 9 + 12)	28189	Foliar	WG	W	3	314 to 395 g/acre (775 to 975 g/ha)	None	7 to 10	12	1	Botrytis grey mould, Monilinia, mummyberry, anthracnose	Sclerotinia	Botrytis cinerea populations with reduced sensitivity to Group 9 fungicides have been observed in some regions of Canada.
	Herbicides													
Broadleaves	<b>Callisto</b> ® (Group 27)	27833	Pre-emergent and post- emergent	SC	L	1	121 mL/acre 0.3 L/ha	Post- emergent: Agral® 90 0.2% v/v	NA	12	60	Pre-emergent: Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard Post-emergent: Nightshade, velvetleaf, redroot pigweed	Common ragweed	
Grasses	Venture L (Group 1)	21209	Post- emergent	EC	Е	1	0.24 to 0.81 L/acre (0.6 to 2 L/ha)	None	NA	12	Fruit: 60 Sprout: 420	Annual grasses	Perennial native grasses	Application rate differs by target.


REI = Re-entry interval PHI = Pre-harvest interval

\*For complete information regarding rates, see product label. Rate information is based on product labels. In the case of ranges, the most commonly used rates are listed. Always refer to the label to confirm crop/crop group registration and application information, as well as buffer zones, PPE, etc., before application. Please check with your packer or buyer when using any Syngenta product on berries potentially destined for export outside of Canada and see the Food Chain ID Regulatory Limits by BCGlobaldatabase at https://bcglobal.bryantchristie.com/db#/pesticides/query for a complete list of MRLs. Should you need additional information on export market MRLs, please consult with Syngenta to receive the most up-to-date information.



## Lowbush blueberry quick reference guide





All information is current at time of publication and is subject to change without notice. For more information, contact our Customer Interaction Centre at 1-87-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

Always read and follow label directions. Miravis® Neo refers to Miravis® Neo 300 SE Fungicide. Agral®, Aprovia®, Bravo®, Callisto®, Inspire Super®, Miravis®, Quilt®, Switch®, Venture® and the Syngenta logo are trademarks of a Syngenta Group Company. Megafol™ is a trademark of VALAGRO S.p.A., a Syngenta Group Company. Allegro® is a trademark of ISK Biosciences Corporation. © 2025 Syngenta

250929

